



Physical activity and risk of endometrial cancer in the Norwegian Women and Cancer (NOWAC) Study

Journal:	<i>International Journal of Cancer</i>
Manuscript ID	IJC-16-2034.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Borch, Kristin; UiT The Arctic University of Norway, Department of Community Medicine Weiderpass, Elisabete ; UiT The Arctic University of Norway, Department of Community Medicine; Karolinska Institutet, Department of Medical Epidemiology and biostatistics; Kreftregisteret, Institute of Population-Based Cancer Research; Samfundet Folkhalsan, Genetic Epidemiology Group Braaten, Tonje; Department of Community Medicine Tromsø, Norway, ISM Jareid, Mie; UiT - The Arctic University of Norway, Department of Community Medicine Gavrilyuk, Oxana; University of Tromsø, Institute of Community Medicine Licaaj, Idlir; Department of Community Medicine,
Key Words:	Endometrial cancer, physical activity, Prospective study, population attributable fraction

SCHOLARONE™
Manuscripts

Physical activity and risk of endometrial cancer in the Norwegian Women and Cancer (NOWAC) Study

Kristin B Borch^{1*}, Elisabete Weiderpass^{1, 2, 3, 4}, Tonje Braaten¹, Mie Jareid¹, Oxana A Gavriluk¹, Ildir Licaj¹

*Cooresponding author: Kristin Benjaminsen Borch, email: Kristin.benjaminsen.borch@uit.no

Running title: Physical activity and endometrial cancer

Word count: Abstract: 250; main text: 3 853

Tables: 5 (Supplementary tables: 1)

Keywords: Endometrial cancer; physical activity; prospective study; population attributable fraction.

Novelty and impact statements:

We found evidence of a dose-response association between physical activity and overall endometrial cancer. The novelty include use of repeated measurements for physical activity and confounders combined with multiple imputation to address attrition, which is a particular problem in observational epidemiology. As a nationally representative cohort our study gave us an unique opportunity to calculate robust population attributable fractions. 22% of endometrial cancer could be avoided if women increase their physical activity level.

Abbreviations:

- BMI – body mass index
- CI – confidence interval
- CUP - Continous Update Project
- PA – physical activity
- PAF – population attributable fractions
- HR – hazard ratio
- RR – relative risk
- NOWAC – The Norwegian Women and Cancer Study
- WCRF/AICR - World Cancer Research Fund/American Institute for Cancer Research

Affiliations:

1. Department of Community Medicine, Faculty of Health Sciences, UiT, The Arctic University of Norway, Tromsø, Norway
2. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
3. Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway
4. Genetic Epidemiology Group, Folkhälsan Research Centre, Samfundet Folkhälsan, Helsinki, Finland

Funding:

This project received financial support from the Norwegian Extra Foundation for Health and Rehabilitation through EXTRA funds.

Conflict of interest: The authors declare that they have no conflict of interest.

Author's contributions

KBB, EW and IL designed the study. IL and KBB performed all statistical analyses, and drafted the manuscript. EW, MJ, and OG critically revised the manuscript. TB participated in the statistical analyses and revised the manuscript critically. All authors read and approved the final manuscript.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

52 **Abstract**

53 Few studies have investigated the association between endometrial cancer and physical
54 activity (PA) using repeated measures of PA and different subtypes of endometrial cancer.
55 We aimed to investigate the association between endometrial cancer and PA level at two
56 points in time in women with different body mass index (BMI) profiles, and to calculate the
57 population attributable fraction (PAF) of endometrial cancer for low PA levels.
58 We included 82,759 women with complete information on PA at baseline in the Norwegian
59 Women and Cancer Study; 52,370 had follow-up information on PA. 687 endometrial cancer
60 cases were identified. Multivariate cox proportional hazard models were used to estimate
61 hazard ratios (HR) and 95% confidence intervals (CI). The PAF indicated the proportion of
62 endometrial cancer that could be avoided in the population if these women had a higher PA
63 level.
64 There was a statistically significant association between low PA levels at baseline and follow-
65 up and endometrial cancer risk, with a dose-response trend (lowest PA level: HR=1.60, 95%
66 CI 1.16-2.20; highest PA level: HR=0.73, 95% CI 0.45-1.16 compared to the median).
67 Analyses that included follow-up measurements yielded similar results. 21.9% (95% CI 7.1-
68 34.3) of endometrial cancers could be avoided if women with low PA levels (≤ 4 in a 1-10
69 degree self reported PA scale) increased their PA levels to 5-10.
70 We found an inverse dose-response association between PA and endometrial cancer,
71 independent of BMI. In this nationally representative cohort, 21.9% of endometrial cancers
72 could potentially be avoided if PA levels increased to higher PA levels.

73

74 Introduction

75 Endometrial cancer is the sixth most common cancer, and the most frequent gynecologic
76 malignancy among women in Norway. In 2014, 727 new cases were diagnosed and 81
77 women died of the disease [1]. Established risk factors for endometrial cancer include use of
78 exogenous estrogens unopposed by progestagens, early menarche (10-12 years of age), late
79 menopause, nulliparity, diabetes mellitus, and obesity. Currently, the majority of endometrial
80 cancer, and about half of the cases in postmenopausal women are attributable to being
81 overweight or obese [2-4]. As the population ages and the prevalence of overweight, obesity,
82 and sedentary lifestyle increase, the incidence of endometrial cancer is also expected to
83 increase, especially in postmenopausal women [5, 6]. Thus primary prevention of endometrial
84 cancer through modifiable lifestyle factors is of potential public health importance.

85 Physical activity (PA) is a modifiable lifestyle factor, which is important in the regulation
86 of hormones and metabolic pathways. It is also associated with weight control, and thus may
87 reduce endometrial cancer risk [7-9]. A pooled analysis of nine cohorts from Europe and the
88 United States included 1.44 million participants and found a 21% reduced risk of endometrial
89 cancer associated with recreational PA [10]. However, within individual studies, results are
90 inconsistent [8]. For example, the Women's Health Study did not find any relationship
91 between recreational PA and walking and endometrial cancer risk [11]. Similarly, there was
92 no significant association between total PA – including occupational, recreational, and
93 household-related PA – and endometrial cancer risk in the European Prospective Investigation
94 into Nutrition and Cancer [12]. The 2007 evaluation by the World Cancer Research
95 Fund/American Institute for Cancer Research (WCRF/AICR) concluded that there is a
96 probable relationship between PA and endometrial cancer despite the variety of PA
97 assessments not allowing for meta-analysis on dose-response [7]. This conclusion was
98 supported in their Continuous Update Project (CUP) report from 2013 [8]. Of four cohort

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

99 studies published after the 2013 CUP report [10, 13-15], three found an inverse association
100 between recreational PA and endometrial cancer risk [10, 15, 14], and one did not [13].

101 Few studies have investigated the association between endometrial cancer and PA using a
102 total and repeated measure of PA and different subtypes of endometrial cancer [12, 16, 17].
103 Endometrial cancer is classified as type I (estrogen dependent), which constitutes the majority
104 of cases (about 80%), and type II (estrogen independent), based on clinical, endocrine, and
105 epidemiological observations. The most common histological subtypes of endometrial cancer
106 are endometrioid carcinoma, serous carcinoma, carcinosarcoma, and clear cell carcinoma
107 [18]. While the association between body mass index (BMI) and endometrial cancer is well
108 established, the relationship between PA and endometrial cancer in women with different
109 body sizes remains unclear.

110 The present study aimed to investigate the association between endometrial cancer and PA
111 level at two points in time in women with different BMI profiles in the Norwegian Women
112 and Cancer (NOWAC) Study, and to calculate the population attributable fraction (PAF) of
113 endometrial cancer for low PA levels.

114
115

116 **Methods**

117 **The NOWAC Study**

118 The NOWAC Study is a nationally representative cohort study that has been described in
119 detail previously [19, 20]. Briefly, random samples of Norwegian women aged 30-70 years
120 were invited to participate during three waves of data collection (1991/92, 1996/97, and
121 2003/04) [20]. More than 172,000 women completed a questionnaire with detailed questions
122 regarding lifestyle, diet, and health, and were enrolled in the study (overall response rate:
123 52.7%). The NOWAC Study was approved by the Regional Committee for Medical Research
124 Ethics and the Norwegian Data Inspectorate, and all participants included in the study gave
125 written informed consent.

126 In this analysis we used information from enrollment questionnaires completed in the
127 period from 1996 to 2004 (baseline), and those with follow-up questionnaires completed 6-8
128 years after enrollment. In total 101,321 women completed questionnaires in these periods and
129 were eligible for inclusion in this study. We excluded women with prevalent cancers other
130 than non-melanoma skin cancer at baseline (n=4,454), those who emigrated or died before the
131 start of follow-up (n=20), those with hysterectomy (n=5,426), and those who had missing
132 information on PA level at baseline (n=8,662). Thus, the final analytical study sample
133 consisted of 82,759 women. Follow-up information on PA level, smoking, weight, and height
134 was available for for 52,370 (63.3%) of these women.

136 **Assessment of PA level and covariates**

137 PA level was assessed in the NOWAC questionnaires on a 10-point scale by the following
138 question: *“By physical activity we mean activity both at work and outside work, at home, as
139 well as training/exercise and other physical activity, such as walking, etc. Please mark the
140 number that best describes your level of physical activity; 1 being very low and 10 being very*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

high". This PA scale has been validated [21] and refers to the total amount of PA across different domains, including recreation, occupation, transportation, and household in one global score. Moderate, but significant Spearman's rank correlation coefficients were found (range: 0.36-0.46; $P<0.001$) between PA level at enrollment and concurrent outcomes from criterion measures of a combined sensor monitoring heart rate and movement. The PA scale appeared valid to rank PA level in Norwegian women, but not to quantify a definite dose of PA [21].

Information on the covariates height, weight, age at menarche, parity, oral contraceptive use, menopausal status, age at menopause, hormone therapy use, years of education, smoking status and alcohol consumption, were obtained from NOWAC questionnaires. The women were considered postmenopausal if they stated that the period had stopped or reported use of hormonal therapy if they were ≥ 53 years. This cut-off point is based on the definition used in the the Million Women Study [22], and has been used by the NOWAC study earlier [23]. Information on height and weight was used to calculate BMI (kg/m^2).

Cancer incidence, death, and emigration

Women diagnosed with a primary, invasive, malignant neoplasm of the endometrium (International Statistical Classification of Diseases, Injuries and Causes of Death Revision 7 codes 172.0 [24]) were identified through linkage to the Cancer Registry of Norway, from which date of diagnosis and morphology (International Classification of Diseases for Oncology, 3rd edition) were also obtained. Based on the morphology, endometrial cancers were categorized into overall endometrial cancer (all subtypes), endometrioid carcinoma, type 1 endometrial cancer (adenocarcinoma NOS, endometrioid, and squamous carcinomas), and other subtypes (non-endometrioid or non-type 1).

Information on date of death or emigration was obtained through linkage to the Norwegian National Population Register.

Statistical methods

Analyses using baseline data only

We used Cox proportional hazard regression models to calculate hazard ratios (HRs) with 95% confidence intervals (CIs) comparing five categories of PA level (1-2, 3-4, 5-6, 7-8, and 9-10). PA level 5-6 was set as the reference group. Follow-up time was defined as the interval between age at baseline and age at cancer diagnosis, death, emigration, or the end of follow-up (31 December 2013), whichever came first.

Departures from the proportional hazards assumption in the Cox models were tested through the inclusion of an interaction variable between categories of PA and underlying time (age). A preliminary analysis of baseline data only was used to select the covariates that were adjusted for in the final models. In the preliminary model, we adjusted for: height (in metres), BMI (normal weight: <25, overweight: 25–29.9, obese: ≥ 30 kg/m²), age at menarche (<12, 13-14, 15+ years), parity (0, 1, 2, 3, 4+) , oral contraceptive use (ever/never), menopausal status (premenopausal, perimenopausal, postmenopausal, hormone therapy use ≥ 53 years), hormone therapy use (ever/never), years of education (≤ 9 , >9-12, ≥ 13 years), smoking status (never, former, current), alcohol consumption (grams per day), and diabetes mellitus (yes/no). The removal of each covariate had to be associated with a change in the regression coefficients of at least 10% in any of the categories of PA level to be included in the final model. To test for linear trend, we used the original, 10-point PA scale, modelled as a continuous variable in the analyses. Interactions (log likelihood test) between PA and the above-mentioned categories of BMI, educational attainment and smoking status were tested. The Wald χ^2 statistic was used to test for heterogeneity between normal weight and obese women.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

190

191

192 Analyses using repeated PA measurements

193 We used the method proposed by Hu et al [25], i.e., baseline data was used until follow-up
194 information became available, death, or emigration, whichever occurred first. Thereafter
195 follow-up information was applied until death, emigration, or the end of the study period,
196 which ever occurred first. In the analysis using repeated PA measurements, we also used
197 follow-up information on BMI and smoking once it became available.

198
199 Complementary analysis - multiple imputation

200 Compared to women who did not drop out of the study, those who dropped out of the study at
201 follow-up (n=30,389 (36.7%)), were more often overweight (31.4% vs 29.2%) or obese
202 (10.3% vs 8.6%), more often reported oral contraceptive use (46.6% vs 43.5%) and hormone
203 therapy use (36.7% vs 31.7%), more often had a history of diabetes mellitus (2.9% vs 1.7%),
204 and had fewer years of education (24.3% vs 23.7%). They were also more often current
205 smokers, but on average they had a similar PA level and alcohol consumption as women who
206 did not drop out of the study. In order to deal with dropouts, we used multiple imputation
207 models [26] and compared the results with those of complete-case analyses. Multiple
208 imputation models were used under the assumption that data was missing at random. To
209 reduce sampling variability, we created 20 replicate datasets from the imputation simulation
210 [27]. We used the outcomes overall endometrial cancer, type 1 endometrial cancer, and
211 endometrioid carcinoma. Nelson-Aalen cumulative hazard estimator was included as a
212 predictor in the imputation models [28]. The estimates from the 20 imputed datasets were
213 combined using Rubin’s rules [29]. All the analyses and multiple imputations were done in
214 STATA version 14.0 (Stata Corp, College Station, TX, USA).

215

216 PAF calculation

217 We calculated the PAF to estimate the proportion of endometrial cancer that could be avoided

218 in the population if women had different PA levels, using the formula: $PAF = Pe * (RRe -$ 219 $1) / [Pe * RR + (1 - Pe)]$, where Pe is the proportion of PA level in the study population and RRe is

220 the RR in the model adjusting for BMI (model 1) and the final baseline multivariable

221 proportional hazards regression model (model 2), including all aforementioned confounders

222 and BMI. We calculated two-sided 95% CIs for the PAFs using the PUNAF Stata module

223 [30]. The PA levels were divided into two levels; levels 1 to 4 were classified as exposed to

224 low PA levels and levels 5-10 as unexposed to low PA levels. The PAF was interpreted as the

225 proportion of overall endometrial cancers that would not occur in the average population if

226 PA levels were between 5 and 10 according to the scale.

227

228

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

The median age at baseline was 51 years. During a mean follow-up time of nearly 13 years, 687 cases of endometrial cancer were identified, with type 1 endometrial cancer (adenocarcinoma not otherwise specified [NOS], endometrioid and squamous carcinoma) accounting for 83.8% of the cases. The age at diagnosis ranged between 42 and 86 years, with a mean of 62 years of age. Nearly half (43%) of the women reported a PA level between 5 and 6 (Table 1), and approximately 74% of the women reported a PA level between 5 and 10. Women with a PA level >5 had a lower BMI, used less hormone therapy, reported less diabetes mellitus, and were more often never smokers compared to women reporting low PA levels.

Multivariable models of the associations between baseline and one follow-up measurement of PA and endometrial cancer risk shows that compared to women with a PA level of 5-6, those with a PA level of 1-2 had a 60% higher risk of overall endometrial cancer (HR=1.60, 95% CI 1.16-2.20). For analyses using repeated PA measurements, the corresponding risks adjusted for BMI and smoking status were similar to those obtained when using baseline data only (HR=1.54, 95% CI 1.01-2.). In analyses using baseline data only and those using repeated PA measurements, BMI and smoking status were negatively associated with the risk of type 1 endometrial cancer and endometrioid carcinomas. Compared to the age-adjusted models, the associations were attenuated in the multivariable adjusted models that included BMI (Table 2).

Interactions between PA and categories of BMI, educational attainment and smoking status were not significant. However, as BMI is a strong risk factor for endometrial cancer, we decided to investigate the association between PA and endometrial cancer risk in normal weight, overweight, and obese women, even though the interaction term was not significant (p=0.49). When analyses were stratified by BMI category the PA levels 7-10 were collapsed,

PA was not associated with overall endometrial cancer among normal-weight women in analyses using baseline data only ($HR_{PA(1-2) \text{ vs } (5-6)}=1.32$, 95% CI 0.71-2.45). The corresponding association in obese participants was $HR_{PA(1-2) \text{ vs } (5-6)}=3.08$ (95% CI 1.76-5.39) ($p_{\text{heterogeneity}}=0.05$) (Table 3).

Using multiple imputation, we found that the estimates at all levels of adjustment and for all endpoints (overall endometrial cancer, type 1 endometrial cancer, and endometrioid carcinoma) were consistent with those obtained from the complete-case analyses (Table 4). PAF calculations showed that if women with a PA level ≤ 4 increased to levels 5-10 in the scale, 21.9% (95% CI: 7.1-34.3) of endometrial cancers could be avoided (Table 5). PAF calculations based on the proportional hazards regression model including BMI yielded a lower proportion (17%, 95% CI: 2.3-29.5) (Table 5). The results did not differ substantially for subtypes of endometrial cancer (Supplemental table 1)

Discussion

In this large Norwegian cohort we found an inverse dose-response association between PA and endometrial cancer overall, type 1 endometrial cancer, and endometrioid carcinoma. The results were consistent when using baseline data on PA and when using repeated PA measurements, as well as when multiple imputation was used. Our findings further suggest that the association between PA and endometrial cancer is independent of BMI, as risk estimates were attenuated, but still significant, when BMI was incorporated in the statistical models. Stratification by BMI category indicated that the risk of low PA on endometrial cancer was statistically significant among obese women only (although the test for interaction was not statistically significant, $p=0.49$). Among women with PA levels ≤ 4 , 21.9% of endometrial cancer could potentially be avoided if these women adopted a PA level between 5-10 in the scale.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Other cohort studies investigating the association between total PA levels – which include domains such as recreation, occupation, transportation, and household – and endometrial cancer are sparse; most studies measured recreational PA only and few studies use repeated measurements of PA. In contrast to our results, the European Prospective Investigation into nutrition and Cancer reported a non-significant trend (p-trend 0.36) for total PA (including the domains recreation, occupation, and household) when comparing active with inactive women (multivariable adjusted model including BMI: HR=0.88, 95% CI 0.61-1.27) [12]. This was similar for the Breast Cancer Detection Demonstration Project study, which measured the total intensity of PA and found no significant associations with endometrial cancer [31]. Findings from the Netherlands Cohort Study on Diet and Cancer showed that total baseline non-occupational PA was inversely associated with endometrial cancer, with a lower risk observed for a PA corresponding to >30 minutes per day [17, 32]. In a Swedish cohort the risk was decreased, although this decrease was not significant [16].

Several studies have reported on recreational PA and endometrial cancer. Nine prospective cohort studies on recreational PA were included in the WCRF/AICR CUP report up to 2012 [8], but due to different measures of PA, the meta-analysis of these studies only looked at the highest vs lowest PA level (RR=0.73, 95% CI 0.58-0.93), and these results were attenuated when the model was adjusted for BMI (RR=0.80, 95% CI 0.69-0.92). A limitation of this meta-analysis was the high heterogeneity ($I^2=75.9\%$) of the individual studies [8]. Indeed, only three of these studies [33, 32, 34] found significant inverse associations, three found no significant association [35, 11, 36], and three [37, 12, 16] found an inverse, but insignificant association between endometrial cancer and recreational PA. Since the publication of the WCRF/AICR CUP report from 2012 we have identified four additional prospective cohort studies on recreational PA and endometrial cancer [10, 13-15]. The Nurses' Health study investigated recreational PA in the past year and found no association

304 between baseline recreational PA and endometrial cancer risk, however, brisk walking time
305 ≥ 3 hours per week was inversely associated with endometrial cancer [15]. In the California
306 Teachers Study cohort, moderate and vigorous recreational PA was associated with a 25%
307 lower endometrial cancer risk [13]. The findings of Land et al [14] are in accordance with our
308 findings, although they studied recreational PA and had a small number of cases in a
309 population of women at high risk for breast cancer. A large pooled analysis of 12 cohorts was
310 recently published and reported a HR of 0.79 (95% CI 0.68-0.92) in the association between
311 recreational PA and endometrial cancer based on 5346 cases. However, the degree of
312 heterogeneity between cohorts was high ($I^2=69\%$)[10].

313 Analysis of occupational PA measured twice during 1974 and 1981 in a Norwegian study
314 showed a significant trend, with a reduced risk for women who were consistently moderately
315 active; however, the association was attenuated in the multivariable model and the trend was
316 no longer significant [36]. The WCRF/AICR CUP report [8] included four cohort studies [16,
317 38, 12, 39], none of which found a significant association between occupational PA and
318 endometrial cancer. In a highest vs lowest occupational PA meta-analysis, the summary RR
319 was 0.79 (95% CI 0.71-0.88), with a high degree of heterogeneity between studies ($I^2=$
320 75.9%) and concluded a probable inverse association between occupational PA and
321 endometrial cancer [8].

322 Overweight and obesity are strong risk factors for endometrial cancer, and studies suggest
323 that the association between PA and endometrial cancer is either mediated or confounded by
324 body weight, which can affect hormone profiles. Therefore, it is important to model the
325 association both with and without adjustment for BMI [15]. In our study, adjustment for BMI
326 in the multivariable analyses attenuated the associations. However, a modest inverse
327 association remained, suggesting that PA is independently associated with endometrial
328 cancer. Simultaneously, the analyses of the different BMI categories showed that the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

association was more pronounced in obese than in normal-weight women. In our data, obesity may confound the association between PA and overall endometrial cancer. Our findings are in accordance with Friberg et al [16]. However, Moore et al adjusted for BMI in the association between recreational PA and endometrial cancer and showed an attenuation of the estimates from a significant towards a non-significant result compared to multivariable models unadjusted for BMI (HR=0.98, 95% CI 0.89-1.09 vs HR=0.79, 95% CI 0.68-0.92) . This was similar to several previous studies [14, 15, 33, 37]. Some studies have found a statistically significant increased endometrial cancer risk in both inactive and active overweight women [11], which correspond to our findings. Others have shown an effect modification, where the inverse relationship was only seen among overweight or obese women [10, 37]. As in our study, several other studies have reported no significant effect modification for BMI [33, 12, 16, 17, 31].

Heterogeneity in different study results may be explained by variations in the methods used to assess PA (self-administered questionnaires, interviews, or use of job titles); PA domains (recreation, occupation, transportation, household); frequency, duration, and intensity of PA; and time periods in life when PA was measured, as well as different statistical methods used in the data analysis [40]. Nevertheless, there is substantial biological evidence to support a potential protective role of PA on endometrial cancer. The mechanisms involved have been hypothesized as affecting endogenous sex hormone levels, insulin-mediated pathways, and maintenance of energy balance [41].

Physical inactivity is considered an important risk factor for different cancers [42-44]. The consistent associations between low PA levels and endometrial cancer risk in our study justify the estimation of PAFs. Our definition of low PA levels was based on self-reported PA using a validated 10-point scale [21]. Our PAF estimation represents the minimum move required from low to higher levels of PA to create a significant change in the incidence of

endometrial cancer (21.9%, 95% CI: 7.1-34.3). However, the definite dose cannot be quantified and our results must be interpreted with caution. In the UK, Parkin found a PAF for endometrial cancer of 3.8% attributable to exercising less than the minimum recommended amount [43]. The proportion related to inadequate PA in the UK in 2002 was 30% for endometrial cancer, however that compared the highest (≥ 60 minutes) and lowest (< 30 minutes) PA levels, which gives a higher reference category than the recommendation of PA [45]. To quantify the PAF requires a realistic population distribution of the exposure of interest, which in our study is PA. We consider our cohort to be a nationally representative Norwegian cohort with a reliable population distribution of PA, and as such it should give a robust estimate. Furthermore, it is valuable to evaluate the impact of different factors in cancer prevention, which is helpful in prioritizing cancer prevention and intervention strategies.

A major strength of our study is its prospective, population-based design, and the use of a high-quality, national cancer registry to identify endometrial cancer cases [20]. The large sample size and representativeness of the Norwegian female population 30 to 70 years of age gives a unique opportunity to calculate robust PAF estimates. The PA scale has been validated [21] and correlated well with all-cause mortality rates [46]. Furthermore, PA level, BMI, and smoking were re-assessed at follow-up. Self-reported BMI has been validated for the NOWAC study, indicating that there was a substantial agreement between self-reported and measured BMI values [47]. There was, however, a small but statistically significant under-reporting of weight, which would affect self-reported BMI; this tendency was largest among overweight and obese women [20, 47]. A survey of the PA levels in the adult population in Norway showed that 34% of women reached the national guidelines for PA [48]. This proportion is higher than in our study. The relatively large number of cases made it possible to investigate subtypes of endometrial cancer, however the proportion of type 2 endometrial

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

cancer was too low to allow for separate analyses. Multiple imputation of missing data, in addition to complete-case analysis, confirmed our results. The PA assessment in our study comprised all areas of PA, not only recreational PA. However, the total self-reported measure of PA cannot differentiate intensity, duration, and frequency of PA, nor the type of PA in our study, and given the self report of PA, measurement errors cannot be ruled out. However, measurement errors would likely lead to a non-differential bias and a potential underestimation of the true effect. The PA assessment used in this study may not apply to women in other countries. Moreover, the potential for residual confounding, in particular by BMI, remains.

Conclusions

Overall, we found an inverse dose-response association between PA and endometrial cancer with similar findings for subtypes of endometrial cancer. This risk was higher in obese women. Also, 21.9% of the endometrial cancer cases could be attributable to low levels of PA, and could potentially be avoided if women attained a higher PA level.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

References

1. Cancer in Norway 2014 - Cancer incidence, mortality, survival and prevalence in Norway. Oslo: Cancer Registry of Norway. 2015.
2. Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D. Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ*. 2007;335(7630):1134. doi:10.1136/bmj.39367.495995.AE.
3. Schmid D, Behrens G, Keimling M, Jochem C, Ricci C, Leitzmann M. A systematic review and meta-analysis of physical activity and endometrial cancer risk. *European Journal of Epidemiology*. 2015;30(5):397-412. doi:10.1007/s10654-015-0017-6.
4. Gil KM, Von Gruening VE. Physical Activity and Gynecologic Cancer Survivorship. In: Courneya KS, Friedenreich CM, editors. *Physical Activity and Cancer*. Berlin Heidelberg: Springer-Verlag 2011.
5. Lacey JV, Jr., Chia VM, Rush BB, Carreon DJ, Richesson DA, Ioffe OB, Ronnett BM, Chatterjee N, Langholz B, Sherman ME, Glass AG. Incidence rates of endometrial hyperplasia, endometrial cancer and hysterectomy from 1980 to 2003 within a large prepaid health plan. *Int J Cancer*. 2012;131(8):1921-9. doi:10.1002/ijc.27457.
6. Amant F, Moerman P, Neven P, Timmerman D, Van Limbergen E, Vergote I. Endometrial cancer. *The Lancet*. 2005;366(9484):491-505.
7. World Cancer Research Fund / American Institute for Cancer Research. *Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global perspective*. Washington DC: AICR; 2007.
8. World Cancer Research Fund/American Institute for Cancer Research. *Continuous Update Project Report. Food, Nutrition, Physical Activity, and the prevention of Endometrial Cancer*. Available at <http://www.dietandcancerreport.org>. 2013.
9. Lindemann K, Eskild A, Vatten LJ, Bray F. Endometrial cancer incidence trends in Norway during 1953–2007 and predictions for 2008–2027. *International Journal of Cancer*. 2010;127(11):2661-8. doi:10.1002/ijc.25267.
10. Moore SC, Lee I, Weiderpass E, Campbell PT, Sampson JN, Kitahara CM, Keadle SK, Arem H, Gonzalez ABd, Hartge P, Adami H-O, Blair CK, Borch KB, Boyd E, Check DP, Fournier A, Freedman ND, Gunter M, Johansson M, Khaw K-T, Linet MS, Orsini N, Park Y, Riboli E, Robien K, Schairer C, Sesso H, Spriggs M, Dusen RV, Wolk A, Matthews CE, Patel AV. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA internal medicine*. 2016;176(6):816-25. doi:10.1001/jamainternmed.2016.1548.
11. Conroy MB, Sattelmair JR, Cook NR, Manson JE, Buring JE, Lee I-M. Physical activity, adiposity, and risk of endometrial cancer. *Cancer Causes & Control*. 2009;20(7):1107-15. doi:10.1007/s10552-009-9313-3.
12. Friedenreich C, Cust A, Lahmann PH, Steindorf K, Boutron-Ruault M-C, Clavel-Chapelon F, Mesrine S, Linseisen J, Rohrmann S, Pischon T, Schulz M, Tjønneland A, Johnsen NF, Overvad K, Mendez M, Arguelles MV, Garcia CM, Larrañaga N, Chirlaque M-D, Ardanaz E, Bingham S, Khaw K-T, Allen N, Key T, Trichopoulos A, Dilis V, Trichopoulos D, Pala V, Palli D, Tumino R, Panico S, Vineis P, Bueno-de-Mesquita HB, Peeters PHM, Monninkhof E, Berglund G, Manjer J, Slimani N, Ferrari P, Kaaks R, Riboli E. Physical activity and risk of endometrial cancer: The European prospective investigation into cancer and nutrition. *International Journal of Cancer*. 2007;121(2):347-55. doi:10.1002/ijc.22676.
13. Dieli-Conwright CM, Ma H, Lacey JV, Henderson KD, Neuhausen S, Horn-Ross PL, Deapen D, Sullivan-Halley J, Bernstein L. Long-term and baseline recreational physical activity and risk of endometrial cancer: the California Teachers Study. *Br J Cancer*. 2013;109(3):761-8. doi:10.1038/bjc.2013.61.
14. Land SR, Liu Q, Wickerham DL, Costantino JP, Ganz PA. Cigarette Smoking, Physical Activity, and Alcohol Consumption as Predictors of Cancer Incidence among Women at High Risk of Breast Cancer in the NSABP P-1 Trial. *Cancer Epidemiology Biomarkers & Prevention*. 2014;23(5):823-32. doi:10.1158/1055-9965.epi-13-1105-t.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

15. Du M, Kraft P, Eliassen AH, Giovannucci E, Hankinson SE, De Vivo I. Physical activity and risk of endometrial adenocarcinoma in the Nurses' Health Study. *International Journal of Cancer*. 2014;134(11):2707-16. doi:10.1002/ijc.28599.

16. Friberg E, Mantzoros CS, Wolk A. Physical Activity and Risk of Endometrial Cancer: A Population-Based Prospective Cohort Study. *Cancer Epidemiology Biomarkers & Prevention*. 2006;15(11):2136-40. doi:10.1158/1055-9965.epi-06-0465.

17. Schouten LJ, Goldbohm RA, van den Brandt PA. Anthropometry, Physical Activity, and Endometrial Cancer Risk: Results From The Netherlands Cohort Study. *Journal of the National Cancer Institute*. 2004;96(21):1635-8. doi:10.1093/jnci/djh291.

18. Murali R, Soslow RA, Weigelt B. Classification of endometrial carcinoma: more than two types. *The Lancet Oncology*. 2014;15(7):e268-e78. doi:[http://dx.doi.org/10.1016/S1470-2045\(13\)70591-6](http://dx.doi.org/10.1016/S1470-2045(13)70591-6).

19. Lund E, Kumle M, Braaten T, Hjartåker A, Bakken K, Eggen A, Gram I. External validity in a population-based national prospective study – the Norwegian Women and Cancer Study (NOWAC). *Cancer Causes and Control*. 2003;14(10):1001-8. doi:10.1023/B:CACO.0000007982.18311.2e.

20. Lund E, Dumeaux V, Braaten T, Hjartåker A, Engeset D, Skeie G, Kumle M. Cohort Profile: The Norwegian Women and Cancer Study (NOWAC) Kvinner og kreft. *Int J Epidemiol*. 2008;37(1):36-41. doi:10.1093/ije/dym137.

21. Borch KB, Ekelund U, Brage S, Lund E. Criterion validity of a 10-category scale for ranking physical activity in Norwegian women. *Int J Behav Nutr Phys Act*. 2012;9(1):2.

22. Collaborators MWS. Breast cancer and hormone-replacement therapy in the Million Women Study. *The Lancet*. 2003;362(9382):419-27. doi:[http://dx.doi.org/10.1016/S0140-6736\(03\)14065-2](http://dx.doi.org/10.1016/S0140-6736(03)14065-2).

23. Waaseth M, Bakken K, Dumeaux V, Olsen K, Rylander C, Figenschau Y, Lund E. Hormone replacement therapy use and plasma levels of sex hormones in the Norwegian Women and Cancer Postgenome Cohort - a cross-sectional analysis. *BMC Women's Health*. 2008;8(1):1.

24. WHO. International Statistical Classification of Diseases and Related Health Problems 10th Revision. <http://apps.who.int/classifications/icd10/browse/2015/en>. 2010. Accessed 27.11.2014.

25. Hu P, Tsiatis AA, Davidian M. Estimating the Parameters in the Cox Model When Covariate Variables are Measured with Error. *Biometrics*. 1998;54(4):1407-19. doi:10.2307/2533667.

26. White IR, Carlin JB. Bias and efficiency of multiple imputation compared with complete-case analysis for missing covariate values. *Statistics in Medicine*. 2010;29(28):2920-31. doi:10.1002/sim.3944.

27. Sterne J, White I, Carlin J, Spratt M, Royston P, Kenward M, Wood A, Carpenter, Jr. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. 2009. doi:10.1136/bmj.b2393.

28. White IR, Royston P. Imputing missing covariate values for the Cox model. *Statistics in Medicine*. 2009;28(15):1982-98. doi:10.1002/sim.3618.

29. Rubin DB. Multiple Imputation After 18+ Years. *Journal of the American Statistical Association*. 1996;91(434):473-89. doi:10.2307/2291635.

30. Benedict N. PUNAF: Stata module to compute population attributable fractions for cohort studies. <https://ideas.repec.org/c/boc/bocode/s457193.html>. 2010. Accessed Accessed 23 June 2016.

31. Colbert LH, Lacey JV, Schairer C, Albert P, Schatzkin A, Albanes D. Physical activity and risk of endometrial cancer in a prospective cohort study (United States). *Cancer Causes & Control*. 2003;14(6):559-67. doi:10.1023/a:1024866827775.

32. Schouten LJ, Goldbohm RA, van den Brandt PA. Anthropometry, Physical Activity, and Endometrial Cancer Risk: Results From The Netherlands Cohort Study. *International Journal of Gynecological Cancer*. 2006;16 (suppl. 2):492.

33. Gierach GL, Chang S-C, Brinton LA, Lacey JV, Hollenbeck AR, Schatzkin A, Leitzmann MF. Physical activity, sedentary behavior, and endometrial cancer risk in the NIH-AARP Diet and Health Study. *International Journal of Cancer*. 2009;124(9):2139-47. doi:10.1002/ijc.24059.

34. Terry P, Baron JA, Weiderpass E, Yuen J, Lichtenstein P, Nyrén O. Lifestyle and endometrial cancer risk: a cohort study from the Swedish twin registry. *International Journal of Cancer*. 1999;82(1):38-42. doi:10.1002/(SICI)1097-0215(19990702)82:1<38::AID-IJC8>3.0.CO;2-Q.
35. Folsom AR, Demissie Z, Harnack L. Glycemic index, glycemic load, and incidence of endometrial cancer: the Iowa Women's Health Study. *Nutrition & Cancer*. 2003;46(2):119-24 6p.
36. Furberg A-S, Thune I. Metabolic abnormalities (hypertension, hyperglycemia and overweight), lifestyle (high energy intake and physical inactivity) and endometrial cancer risk in a Norwegian cohort. *International Journal of Cancer*. 2003;104(6):669-76. doi:10.1002/ijc.10974.
37. Patel AV, Feigelson HS, Talbot JT, McCullough ML, Rodriguez C, Patel RC, Thun MJ, Calle EE. The role of body weight in the relationship between physical activity and endometrial cancer: Results from a large cohort of US women. *International Journal of Cancer*. 2008;123(8):1877-82. doi:10.1002/ijc.23716.
38. Moradi T, Nyrén O, Bergström R, Gridley G, Linet M, Wolk A, Dosemeci M, Adami H-O. Risk for endometrial cancer in relation to occupational physical activity: A nationwide cohort study in Sweden. *International Journal of Cancer*. 1998;76(5):665-70. doi:10.1002/(SICI)1097-0215(19980529)76:5<665::AID-IJC9>3.0.CO;2-O.
39. Weiderpass E, Pukkala E, Vasama-Neuvonen K, Kauppinen T, Vainio H, Paakkulainen H, Boffetta P, Partanen T. Occupational exposures and cancers of the endometrium and cervix uteri in Finland. *American Journal of Industrial Medicine*. 2001;39(6):572-80. doi:10.1002/ajim.1056.
40. Friedenreich CM. Physical Activity and Cancer Prevention: From Observational to Intervention Research. *Cancer Epidemiology Biomarkers & Prevention*. 2001;10(4):287-301.
41. Cust AE. Physical Activity and Gynecologic Cancer Prevention. *Physical Activity and Cancer*. In: Courneya KS, Friedenreich CM, editors. *Recent Results in Cancer Research*. Berlin Heidelberg: Springer; 2011. p. 159-85.
42. Mathers C, Stevens GA, Mascarenhas M. Global Health Risks: Mortality burden of disease attributable to selected major risk. Switzerland: WHO2009. Report No.: 9789241563871.
43. Parkin DM. 9. Cancers attributable to inadequate physical exercise in the UK in 2010. *Br J Cancer*. 2011;105(S2):S38-S41.
44. Olsen CM, Wilson LF, Nagle CM, Kendall BJ, Bain CJ, Pandeya N, Webb PM, Whiteman DC. Cancers in Australia in 2010 attributable to insufficient physical activity. *Australian and New Zealand Journal of Public Health*. 2015;39(5):458-63. doi:10.1111/1753-6405.12469.
45. World Cancer Research Fund/American Institute for Cancer Research. Policy and Action for Cancer Prevention. Food, Nutrition and Physical Activity: A Global Perspective. AICR: Washington, DC. 2009.
46. Borch KB, Braaten T, Lund E, Weiderpass E. Physical activity and mortality among Norwegian women - the Norwegian Women and Cancer Study. *Clin Epidemiol*. 2011;3(1):229-35. doi:10.1186/1479-5868-9-2.
47. Skeie G, Mode N, Henningsen M, Borch KB. Validity of self-reported body mass index among middle-aged participants in the Norwegian Women and Cancer study. *Clinical Epidemiology*. 2015;7:313-23. doi:10.2147/CLEP.S83839.
48. Hansen BH, Anderssen SA, Steene-Johannessen J, Ekelund U, Nilsen AK, Andersen ID, Dalene KE, Kolle E. Fysisk aktivitet og sedat tid blant voksne og eldre i Norge-Nasjonalt kartlegging 2014-2015. Helsedirektoratet, Oslo2015.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **Physical activity and risk of endometrial cancer in the Norwegian Women**
2 **and Cancer (NOWAC) Study**

3 Kristin B Borch^{1*}, Elisabete Weiderpass^{1, 2, 3, 4}, Tonje Braaten¹, Mie Jareid¹, Oxana A
4 Gavrilyuk¹, Ildir Licaj¹

5 *Cooresponding author: Kristin Benjaminsen Borch, email:
6 Kristin.benjaminsen.borch@uit.no

7 **Running title: Physical activity and endometrial cancer**

8 **Word count:** Abstract: 250; main text: 3 853+9

9 **Tables:** 35+ (supplementary tables: 1-figures:-+)

10 **Keywords:** Endometrial cancer; physical activity; prospective study; population attributable
11 fraction.

12

13 **Novelty and impact statements:**

14 We found evidence of a dose-response association between physical activity and overall
15 endometrial cancer. The novelty include use of repeated measurements for physical activity
16 and confounders combined with multiple imputation to address attrition, which is a particular
17 problem in observational epidemiology. As a nationally representative cohort our study gave
18 us an unique opportunity to calculate robust population attributable fractions. 34+22% of
19 endometrial cancer could be avoided if women increase their physical activity to the
20 recommended level.

21 **Abbreviations:**

22 BMI – body mass index
23 CI – confidence interval
24 CUP - Continous Update Project
25 PA – physical activity
26 PAF – population attributable fractions
27 HR – hazard ratio
28 RR – relative risk
29 NOWAC – The Norwegian Women and Cancer Study
30 WCRF/AICR - World Cancer Research Fund/American Institute for Cancer Research

31

Affiliations:

1. Department of Community Medicine, Faculty of Health Sciences, UiT, The Arctic University of Norway, Tromsø, Norway
2. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
3. Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway
4. Genetic Epidemiology Group, Folkhälsan Research Centre, Samfundet Folkhälsan, Helsinki, Finland

Funding:

This project received financial support from the Norwegian Extra Foundation for Health and Rehabilitation through EXTRA funds.

Conflict of interest: The authors declare that they have no conflict of interest.

Author's contributions

KBB, EW and IL designed the study. IL and KBB performed all statistical analyses, and drafted the manuscript. EW, MJ, and OG critically revised the manuscript. TB participated in the statistical analyses and revised the manuscript critically. All authors read and approved the final manuscript.

1
2
3
4
5
6
7 **Abstract**
8
9 Few studies have investigated the association between endometrial cancer and physical
10 activity (PA) using repeated measures of PA and different subtypes of endometrial cancer.
11
12 We aimed to investigate the association between endometrial cancer and PA level at two
13 points in time in women with different body mass index (BMI) profiles, and to calculate the
14 population attributable fraction (PAF) of endometrial cancer for low PA levels.
15
16 We included 82,759 women with complete information on PA at baseline in the Norwegian
17 Women and Cancer Study; 52,370 had follow-up information on PA. 687 endometrial cancer
18 cases were identified. Multivariate cox proportional hazard models were used to estimate
19 hazard ratios (HR) and 95% confidence intervals (CI). The PAF indicated the proportion of
20 endometrial cancer that could be avoided in the population if these women had a higher PA
21 level, ~~corresponding to ≥ 150 minutes of moderate/vigorous PA per week.~~
22
23 There was a statistically significant association between low PA levels at baseline and follow-
24 up and endometrial cancer risk, with a dose-response trend (lowest PA level: HR=1.~~8560~~,
25 95% CI 1.~~1636-2.2053~~; highest PA level: HR=0.7~~43~~, 95% CI 0.45-1.1~~26~~ compared to the
26 median). ~~The Analyses that included including follow-up measurements yielded similar~~
27 ~~results. so associations were attenuated after adjustment for BMI, but remained significant.~~
28
29 ~~231.94% (95% CI 7.16-34.349) of endometrial cancers could be avoided if women with low~~
30 ~~PA levels (≤ 4 in a 1-10 degree self reported PA scale) increased their PA levels to 5-10 at~~
31 ~~least 150 minutes/week.~~
32
33 We found an inverse dose-response association between PA and endometrial cancer,
34 independent of BMI. ~~In this nationally representative cohort, 21.9% One of four~~ endometrial
35 cancers could potentially be avoided if PA levels increased to higher PA levels (5-10 in the
36 scale) ~~150 minutes per week.~~
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

78 Introduction

79 Endometrial cancer is the sixth most common cancer, and the most frequent gynecologic
80 malignancy among women in Norway. In 2014, 727 new cases were diagnosed and 81
81 women died of the disease [1]. Established risk factors for endometrial cancer include use of
82 exogenous estrogens unopposed by progestagens, early menarche (10-12 years of age), late
83 menopause, nulliparity, diabetes mellitus, and obesity. Currently, the majority of endometrial
84 cancer, and about half of the cases in postmenopausal women are attributable to being
85 overweight or obese [2-4]. As the population ages and the prevalence of overweight, obesity,
86 and sedentary lifestyle increase, the incidence of endometrial cancer is also expected to
87 increase, especially in postmenopausal women [5, 6]. Thus primary prevention of endometrial
88 cancer through modifiable lifestyle factors is of potential public health importance.

89 Physical activity (PA) is a modifiable lifestyle factor, which is important in the regulation
90 of hormones and metabolic pathways. It is also associated with weight control, and thus may
91 reduce endometrial cancer risk [7-9]. A pooled analysis of nine cohorts from Europe and the
92 United States included 1.44 million participants and found a 21% reduced risk of endometrial
93 cancer associated with recreational PA [10]. However, within individual studies, results are
94 inconsistent [8]. For example, the Women's Health Study did not find any relationship
95 between recreational PA and walking and endometrial cancer risk [11]. Similarly, there was
96 no significant association between total PA – including occupational, recreational, and
97 household-related PA – and endometrial cancer risk in the European Prospective Investigation
98 into Nutrition and Cancer [12]. The 2007 evaluation by the World Cancer Research
99 Fund/American Institute for Cancer Research (WCRF/AICR) concluded that there is a
100 probable relationship between PA and endometrial cancer despite the variety of PA
101 assessments not allowing for meta-analysis on dose-response [7]. This conclusion was
102 supported in their Continuous Update Project (CUP) report from 2013 [8]. Of four cohort

1
2
3
4
5
6
7 103 studies published after the 2013 CUP report [10, 13-15], three found an inverse association
8
9 104 between recreational PA and endometrial cancer risk [10, 15, 14], and one did not [13].
10
11 105 Few studies have investigated the association between endometrial cancer and PA using a
12
13 106 total and repeated measure of PA and different subtypes of endometrial cancer [12, 16, 17].
14
15 107 Endometrial cancer is classified as type I (estrogen dependent), which constitutes the majority
16
17 108 of cases (about 80%), and type II (estrogen independent), based on clinical, endocrine, and
18
19 109 epidemiological observations. The most common histological subtypes of endometrial cancer
20
21 110 are endometrioid carcinoma, serous carcinoma, carcinosarcoma, and clear cell carcinoma
22
23 111 [18]. While the association between body mass index (BMI) and endometrial cancer is well
24
25 112 established, the relationship between PA and endometrial cancer in women with different
26
27 113 body sizes remains unclear.
28
29 114 The present study aimed to investigate the association between endometrial cancer and PA
30
31 115 level at two points in time in women with different BMI profiles in the Norwegian Women
32
33 116 and Cancer (NOWAC) Study, and to calculate the population attributable fraction (PAF) of
34
35 117 endometrial cancer for low PA levels.
36
37 118
38 119

120 **Methods**

121 **The NOWAC Study**

122 The NOWAC Study is a nationally representative cohort study that has been described in
123 detail previously [19, 20]. Briefly, random samples of Norwegian women aged 30-70 years
124 were invited to participate during three waves of data collection (1991/92, 1996/97, and
125 2003/04) [20]. More than 172,000 women completed a questionnaire with detailed questions
126 regarding lifestyle, diet, and health, and were enrolled in the study (overall response rate:
127 52.7%). The NOWAC Study was approved by the Regional Committee for Medical Research
128 Ethics and the Norwegian Data Inspectorate, and all participants included in the study gave
129 written informed consent.

130 In this analysis we used information from enrollment questionnaires completed in the
131 period from 1996 to 2004 (baseline), and those with follow-up questionnaires completed 6-8
132 years after enrollment. In total 101,321 women completed questionnaires in these periods and
133 were eligible for inclusion in this study. We excluded women with prevalent cancers other
134 than non-melanoma skin cancer at baseline (n=4,454), those who emigrated or died before the
135 start of follow-up (n=20), those with hysterectomy (n=5,426), and those who had missing
136 information on PA level at baseline (n=8,662). Thus, the final analytical study sample
137 consisted of 82,759 women. Follow-up information on PA level, smoking, weight, and height
138 was available for for 52,370 (63.3%) of these women.

140 **Assessment of PA level and covariates**

141 PA level was assessed in the NOWAC questionnaires on a 10-point scale by the following
142 question: *“By physical activity we mean activity both at work and outside work, at home, as
143 well as training/exercise and other physical activity, such as walking, etc. Please mark the
144 number that best describes your level of physical activity; 1 being very low and 10 being very*

1
2
3
4
5
6
7 145 *high*". This PA scale has been validated [21] and refers to the total amount of PA across
8
9 146 different domains, including recreation, occupation, transportation, and household in one
10
11 147 global score. Moderate, but significant Spearman's rank correlation coefficients were found
12
13 148 (range: 0.36-0.46; $P<0.001$) between PA level at enrollment and concurrent outcomes from
14
15 149 criterion measures of a combined sensor monitoring heart rate and movement. ~~This~~
16
17 150 ~~corresponded to mean values of 0.8 (very low) and 3.4 hours per day (very high) of~~
18
19 151 ~~moderate/vigorous PA, respectively, with a linear increase (P for trend <0.001).~~The PA scale
20
21 152 appeared valid to rank PA level in Norwegian women, but not to quantify a definite dose of
22
23 153 PA [21].
24
25 154 Information on the covariates height, weight, age at menarche, parity, oral contraceptive
26
27 155 use, menopausal status, age at menopause, hormone therapy use, years of education, smoking
28
29 156 status ~~and~~, alcohol consumption, were obtained from NOWAC questionnaires. The women
30
31 157 were as considered postmenopausal if they stated that the period had stopped or had a
32
33 158 hysterectomy (excluded from the study) or reported use of hormonal therapy if they were
34
35 159 >53 years. This cut-off point is based on the definition used in the the Million Women Study
36
37 160 [22], and has been used by the NOWAC study earlier [23]. Information on height and weight
38
39 161 was used to calculate BMI (kg/m^2).
40
41 162

41 163 **Cancer incidence, death, and emigration**

42
43 164 Women diagnosed with a primary, invasive, malignant neoplasm of the endometrium
44
45 165 (International Statistical Classification of Diseases, Injuries and Causes of Death Revisions 7
46
47 166 ~~and 10~~ codes 172.0 ~~and C 54.1, respectively~~ [24]) were identified through linkage to the
48
49 167 Cancer Registry of Norway, from which date of diagnosis and morphology (International
50
51 168 Classification of Diseases for Oncology, 3rd edition) were also obtained. Based on the
52
53 169 morphology, endometrial cancers were categorized into overall endometrial cancer (all
54
55
56
57
58
59
60

subtypes), endometrioid carcinoma, type 1 endometrial cancer (adenocarcinoma NOS, endometrioid, and squamous carcinomas), and other subtypes (non-endometrioid or non-type 1).

Information on date of death or emigration was obtained through linkage to the Norwegian National Population Register.

Statistical methods

Analyses using baseline data only

We used Cox proportional hazard regression models to calculate hazard ratios (HRs) with 95% confidence intervals (CIs) comparing five categories of PA level (1-2, 3-4, 5-6, 7-8, and 9-10). PA level 5-6 was set as the reference group. Follow-up time was defined as the interval between age at baseline and age at cancer diagnosis, death, emigration, or the end of follow-up (31 December 2013), whichever came first.

Departures from the proportional hazards assumption in the Cox models were tested through the inclusion of an interaction variable between categories of PA and underlying time (age). A preliminary analysis of baseline data only was used to select the covariates that were adjusted for in the final models. In the preliminary model, we adjusted for: height (in metres), BMI (normal weight: <25, overweight: 25–29.9, obese: ≥ 30 kg/m²), age at menarche (<12, 13-14, 15+ years), parity (0, 1, 2, 3, 4+) , oral contraceptive use (ever/never), menopausal status (premenopausal, perimenopausal, postmenopausal, hormone therapy use ≥ 53 years), hormone therapy use (ever/never), years of education (≤ 9 , >9-12, ≥ 13 years), smoking status (never, former, current), alcohol consumption (grams per day), and diabetes mellitus (yes/no).

The removal of each covariate had to be associated with a change in the regression coefficients of at least 10% in any of the categories of PA level to be included in the final model. To test for linear trend, we used the original, 10-point PA scale, modelled as a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

continuous variable in the analyses. Interactions (log likelihood test) between PA and the above-mentioned categories of BMI, educational attainment and smoking status were tested. The Wald χ^2 statistic was used to test for heterogeneity between normal weight and obese women, ~~not significant~~. As BMI is a strong risk factor for endometrial cancer, we decided to investigate the association between PA and endometrial cancer risk in normal weight, overweight, and obese women, even though the interaction term was not significant ($p=0.49$). The Wald χ^2 statistic was used to test for heterogeneity between normal weight and obese women.

Analyses using repeated PA measurements

We used the method proposed by Hu et al [25], i.e., baseline data was used until follow-up information became available, death, or emigration, whichever occurred first. Thereafter follow-up information was applied until death, emigration, or the end of the study period, which ever occurred first. In the analysis using repeated PA measurements, we also used follow-up information on BMI and smoking once it became available.

Complementary analysis - multiple imputation

Compared to women who did not drop out of the study, those who dropped out of the study at follow-up ($n=30,389$ (36.7%)), were more often overweight (31.4% vs 29.2%) or obese (10.3% vs 8.6%), more often reported oral contraceptive use (46.6% vs 43.5%) and hormone therapy use (36.7% vs 31.7%), more often had a history of diabetes mellitus (2.9% vs 1.7%), and had fewer years of education (24.3% vs 23.7%). They were also more often current smokers, but on average they had a similar PA level and alcohol consumption as women who did not drop out of the study. In order to deal with dropouts, we used multiple imputation models [26] and compared the results with those of complete-case analyses. Multiple

imputation models were used under the assumption that data was missing at random. To reduce sampling variability, we created 20 replicate datasets from the imputation simulation [27]. We used the outcomes overall endometrial cancer, type 1 endometrial cancer, and endometrioid carcinoma. Nelson-Aalen cumulative hazard estimator was included as a predictor in the imputation models [28]. The estimates from the 20 imputed datasets were combined using Rubin's rules [29]. All the analyses and multiple imputations were done in STATA version 14.0 (Stata Corp, College Station, TX, USA).

PAF calculation

We calculated the PAF to estimate the proportion of endometrial cancer that could be avoided in the population if women had different PA levels, using the formula: $PAF = Pe * (RRe - 1) / [Pe * RR + (1 - Pe)]$, where Pe is the proportion of PA level in the study population and RRe is the RR in the model adjusting for BMI (model 1) and the final baseline multivariable proportional hazards regression model (model 2), including all aforementioned confounders and BMI. We calculated two-sided 95% CIs for the PAFs using the PUNAF Stata module [30]. The PA levels were divided into two levels: levels 1 to 4 were classified as exposed to low PA levels and levels 5-10 as unexposed to low PA levels. The PAF was interpreted as the proportion of overall endometrial cancers that would not occur in the average population if PA levels ~~was~~ were between 5 and 10 according to the scale. ≥ 8 , corresponding to 150 minutes per week according to the validation of PA in NOWAC, assuming that the distribution of the adjustment variables remained unchanged.

1
2
3
4
5
6
7 243 **Results**
8
9 244 The median age at baseline was 51 years. During a mean follow-up time of nearly 13 years,
10 245 687 cases of endometrial cancer were identified, with type 1 endometrial cancer
11
12 246 (adenocarcinoma not otherwise specified [NOS], endometrioid and squamous carcinoma)
13
14 247 accounting for 83.8% of the cases. The age at diagnosis ranged between 42 and 86 years, with
15
16 248 a mean of 62 years of age. Nearly half (43%) of the women reported a PA level between 5
17
18 249 and 6 (Table 1), and collapsing the upper part of the scale showed that approximately 74%
19
20 250 of the women reported a PA level between 5 and 10, with level ≥ 8 comprised 15% of the
21
22 251 participants. Women with a PA level >5 had a lower BMI, used less hormone therapy,
23
24 252 reported less diabetes mellitus, and were more often never smokers compared to women
25
26 253 reporting low PA levels. (Table 1).
27
28 254 Multivariable models of the associations between baseline and one follow-up
29
30 255 measurement of PA and endometrial cancer risk shows that compared to women with a PA
31
32 256 level of 5-6, those with a PA level of 1-2 had a 60% higher risk of overall endometrial cancer
33
34 257 (HR=1.60, 95% CI 1.16-2.20). Removing BMI from the models increased these estimates
35
36 258 (HR=1.85, 95% CI 1.36-2.36). For analyses using repeated PA measurements, the
37
38 259 corresponding risks adjusted for BMI and smoking status were similar to those obtained when
39
40 260 using baseline data only (HR with BMI adjustment =1.54, 95% CI 1.01-2.17 and HR without
41
42 261 BMI adjustment =1.80, 95% CI 1.19-2.72). In analyses using baseline data only and those
43
44 262 using repeated PA measurements, BMI and smoking status were negatively associated with the
45
46 263 risk of type 1 endometrial cancer and endometrioid carcinomas. Compared to the age-
47
48 264 adjusted models, the associations were attenuated in the multivariable adjusted models that
49
50 265 included BMI (Table 2).
51
52 266 Interactions between PA and categories of BMI, educational attainment and smoking
53
54 267 status were not significant. However, as BMI is a strong risk factor for endometrial cancer,
55
56
57
58
59
60

we decided to investigate the association between PA and endometrial cancer risk in normal weight, overweight, and obese women, even though the interaction term was not significant ($p=0.49$). When analyses were stratified by BMI category the PA levels 7-10 were collapsed, PA was not associated with overall endometrial cancer among normal-weight women in analyses using baseline data only ($HR_{PA(1-2) \text{ vs } (5-6)}=1.32$, 95% CI 0.71-2.45). The corresponding association in obese participants was $HR_{PA(1-2) \text{ vs } (5-6)}=3.08$ (95% CI 1.76-5.39) ($p_{\text{heterogeneity}}=0.05$) (Table 3).

Using multiple imputation, we found that the estimates at all levels of adjustment and for all endpoints (overall endometrial cancer, type 1 endometrial cancer, and endometrioid carcinoma) were consistent with those obtained from the complete-case analyses (Table 4). (Supplemental Table 1). PAF calculations showed that if women with a PA level ≤ 4 increased to levels 5-10 in the scale that level to ≥ 8 , 21.934% (95% CI: 7.16-34.349) of endometrial cancers could be avoided (Table 5). PAF calculations based on the proportional hazards regression model including BMI yielded a lower proportion (17%, 95% CI: 2.3-29.5) (Table 5). The results did not differ substantially for subtypes of endometrial cancer (Supplemental table 1) (more than 1 in 4) (Supplementary Figure 1).

Discussion

In this large Norwegian cohort we found an inverse dose-response association between PA and endometrial cancer overall, type 1 endometrial cancer, and endometrioid carcinoma. The results were consistent when using baseline data on PA and when using repeated PA measurements, as well as when multiple imputation was used. Our findings further suggest that the association between PA and endometrial cancer is independent of BMI, as risk estimates were attenuated, but still significant, when BMI was incorporated in the statistical models. Stratification by BMI category indicated that the risk of low PA on endometrial cancer was statistically significant among obese women only (although the test for interaction

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

was not statistically significant, $p=0.49$). Among women with PA levels ≤ 4 , 21.93% of endometrial cancer could potentially be avoided if these women adopted a PA level between 5-10 in the scale ≥ 8 , which corresponds to approximately 150 minutes of moderate/vigorous PA per week.

Other cohort studies investigating the association between total PA levels – which include domains such as recreation, occupation, transportation, and household – and endometrial cancer are sparse; most studies measured recreational PA only and few studies use repeated measurements of PA. In contrast to our results, the European Prospective Investigation into nutrition and Cancer reported a non-significant trend (p-trend 0.36) for total PA (including the domains recreation, occupation, and household) when comparing active with inactive women (multivariable adjusted model including BMI: HR=0.88, 95% CI 0.61-1.27) [12]. This was similar for the Breast Cancer Detection Demonstration Project study, which measured the total intensity of PA and found no significant associations with endometrial cancer [31]. Findings from the Netherlands Cohort Study on Diet and Cancer showed that total baseline non-occupational PA was inversely associated with endometrial cancer, with a lower risk observed for a PA corresponding to >30 minutes per day [17, 32]. In a Swedish cohort the risk was decreased, although this decrease was not significant [16].

Several studies have reported on recreational PA and endometrial cancer. Nine prospective cohort studies on recreational PA were included in the WCRF/AICR CUP report up to 2012 [8], but due to different measures of PA, the meta-analysis of these studies only looked at the highest vs lowest PA level (RR=0.73, 95% CI 0.58-0.93), and these results were attenuated when the model was adjusted for BMI (RR=0.80, 95% CI 0.69-0.92). A limitation of this meta-analysis was the high heterogeneity ($I^2=75.9\%$) of the individual studies [8]. Indeed, only three of these studies [33, 32, 34] found significant inverse associations, three found no significant association [35, 11, 36], and three [37, 12, 16] found an inverse, but

insignificant association between endometrial cancer and recreational PA. Since the publication of the WCRF/AICR CUP report from 2012 we have identified four additional prospective cohort studies on recreational PA and endometrial cancer [10, 13-15]. The Nurses' Health study investigated recreational PA in the past year and found no association between baseline recreational PA and endometrial cancer risk, however, brisk walking time ≥ 3 hours per week was inversely associated with endometrial cancer [15]. In the California Teachers Study cohort, moderate and vigorous recreational PA was associated with a 25% lower endometrial cancer risk [13]. The findings of Land et al [14] are in accordance with our findings, although they studied recreational PA and had a small number of cases in a population of women at high risk for breast cancer. A large pooled analysis of 12 cohorts was recently published and reported a HR of 0.79 (95% CI 0.68-0.92) in the association between recreational PA and endometrial cancer based on 5346 cases. However, the degree of heterogeneity between cohorts was high ($I^2=69\%$)[10].

Analysis of occupational PA measured twice during 1974 and 1981 in a Norwegian study showed a significant trend, with a reduced risk for women who were consistently moderately active; however, the association was attenuated in the multivariable model and the trend was no longer significant [36]. The WCRF/AICR CUP report [8] included four cohort studies [16, 38, 12, 39], none of which found a significant association between occupational PA and endometrial cancer. In a highest vs lowest occupational PA meta-analysis, the summary RR was 0.79 (95% CI 0.71-0.88), with a high degree of heterogeneity between studies ($I^2=75.9\%$) and concluded a probable inverse association between occupational PA and endometrial cancer [8].

Overweight and obesity are strong risk factors for endometrial cancer, and studies suggest that the association between PA and endometrial cancer is either mediated or confounded by body weight, which can affect hormone profiles. Therefore, it is important to model the

1
2
3
4
5
6
7 344 association both with and without adjustment for BMI [15]. In our study, adjustment for BMI
8
9 345 in the multivariable analyses attenuated the associations. However, a modest inverse
10
11 346 association remained, suggesting that PA is independently associated with endometrial
12
13 347 cancer. Simultaneously, the analyses of the different BMI categories showed that the
14
15 348 association was more pronounced in obese than in normal-weight women. In our data, obesity
16
17 349 may confound the association between PA and overall endometrial cancer. Our findings are in
18
19 350 accordance with Friberg et al [16]. However, Moore et al adjusted for BMI in the association
20
21 351 between recreational PA and endometrial cancer and showed an attenuation of the estimates
22
23 352 from a significant towards a non-significant result compared to multivariable models
24
25 353 unadjusted for BMI (HR=0.98, 95% CI 0.89-1.09 vs HR=0.79, 95% CI 0.68-0.92) . This was
26
27 354 similar to several previous studies [14, 15, 33, 37]. Some studies have found a statistically
28
29 355 significant increased endometrial cancer risk in both inactive and active overweight women
30
31 356 [11], which correspond to our findings. Others have shown an effect modification, where the
32
33 357 inverse relationship was only seen among overweight or obese women [10, 37]. As in our
34
35 358 study, several other studies have reported no significant effect modification for BMI [33, 12,
36
37 359 16, 17, 31].

38
39 360 Heterogeneity in different study results may be explained by variations in the methods
40
41 361 used to assess PA (self-administered questionnaires, interviews, or use of job titles); PA
42
43 362 domains (recreation, occupation, transportation, household); frequency, duration, and
44
45 363 intensity of PA; and time periods in life when PA was measured, as well as different statistical
46
47 364 methods used in the data analysis [40]. Nevertheless, there is substantial biological evidence
48
49 365 to support a potential protective role of PA on endometrial cancer. The mechanisms involved
50
51 366 have been hypothesized as affecting endogenous sex hormone levels, insulin-mediated
52
53
54
55
56
57
58
59
60 367 pathways, and maintenance of energy balance [41].

Physical inactivity is considered an important risk factor for different cancers [42-44]. The consistent associations between low PA levels and endometrial cancer risk in our study justify the estimation of PAFs. Our definition of low PA levels was based on self-reported PA using a validated 10-point scale [21]. Our PAF estimation represents the minimum move required from low to higher levels of PA to create a significant change in the incidence of endometrial cancer (21.934%, 95% CI: 7.1-34.3). However, the definite dose cannot be quantified and our results must be interpreted with caution, which corresponded to an increase in PA to 150 minutes of moderate/vigorous PA per week. This amount is in accordance with World Health Organization Global Recommendations on PA for Health. In the UK, Parkin found a PAF for endometrial cancer of 3.8% attributable to exercising less than the minimum recommended amount [43]. The proportion related to inadequate PA in the UK in 2002 was 30% for endometrial cancer, however that compared the highest (≥ 60 minutes) and lowest (< 30 minutes) PA levels, which gives a higher reference category than the recommendation of PA [45]. To quantify the PAF requires a realistic population distribution of the exposure of interest, which in our study is PA. We consider our cohort to be a nationally representative Norwegian cohort with a reliable population distribution of PA, and as such it should give a more-robust estimate. Furthermore, it is valuable to evaluate the impact of different factors in cancer prevention, which is helpful in prioritizing cancer prevention and intervention strategies.

A major strength of our study is its prospective, population-based design, and the use of a high-quality, nationally representative cancer registry system to identify endometrial cancer cases [20]. The large sample size and representativeness of the Norwegian female population 30 to 70 years of age gives a unique opportunity to calculate robust PAF estimates. The PA scale has been validated [21] and correlated well with all-cause mortality rates [46]. Furthermore, PA level, BMI, and smoking were re-assessed at follow-up. Self-reported BMI

1
2
3
4
5
6
7 393 has been validated for the NOWAC study, indicating that there was a substantial agreement
8
9 394 between self-reported and measured BMI values [47]. There was, however, a small but
10
11 395 statistically significant under-reporting of weight, which would affect self-reported BMI; this
12
13 396 tendency was largest among overweight and obese women [20, 47] . A survey of the PA
14
15 397 levels in the adult population in Norway showed that 34% of women reached the national
16
17 398 guidelines for PA [48]. This proportion is higher than in our study. The relatively large
18
19 399 number of cases made it possible to investigate subtypes of endometrial cancer, however the
20
21 400 proportion of type 2 endometrial cancer was too low to allow for separate analyses. Multiple
22
23 401 imputation of missing data, in addition to complete-case analysis, confirmed our results. The
24
25 402 PA assessment in ~~our~~^{this} study comprised all areas of PA, not only recreational PA. However,
26
27 403 the total self-reported measure of PA cannot differentiate intensity, duration, and frequency of
28
29 404 PA, nor the type of PA in our study, and given the self report of PA, measurement errors
30
31 405 cannot be ruled out. However, measurement errors would likely lead to a non-differential bias
32
33 406 and a potential underestimation of the true effect. The PA assessment used in this study may
34
35 407 not apply to women in other countries. Moreover, the potential for residual confounding, in
36
37 408 particular by BMI, remains.

37 409
38
39 410 **Conclusions**
40
41 411 Overall, we found an inverse dose-response association between PA and endometrial cancer
42
43 412 with similar findings for subtypes of endometrial cancer. This risk was higher in obese
44
45 413 women. Also, ~~21.934~~^{21.93}% of the endometrial cancer cases ~~could be were~~ attributable to low
46
47 414 levels of PA, and ~~more than 1 in 4 cases~~ could potentially be avoided if women attained a
48
49 415 higher PA level ~~corresponding to 150 minutes per week or more.~~

50 416
51
52
53 417
54
55
56
57
58
59
60

1
2
3
4
5
6
7 418 **Ethical approval:** All procedures performed in studies involving human participants were in
8
9 419 accordance with the ethical standards of the institutional and/or national research committee
10
11 420 and with the 1964 Helsinki declaration and its later amendments or comparable ethical
12
13 421 standards.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Cancer in Norway 2014 - Cancer incidence, mortality, survival and prevalence in Norway. Oslo: Cancer Registry of Norway. 2015.

2. Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D. Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ*. 2007;335(7630):1134. doi:10.1136/bmj.39367.495995.AE.

3. Schmid D, Behrens G, Keimling M, Jochem C, Ricci C, Leitzmann M. A systematic review and meta-analysis of physical activity and endometrial cancer risk. *European Journal of Epidemiology*. 2015;30(5):397-412. doi:10.1007/s10654-015-0017-6.

4. Gil KM, Von Grueningen VE. Physical Activity and Gynecologic Cancer Survivorship. In: Courneya KS, Friedenreich CM, editors. *Physical Activity and Cancer*. Berlin Heidelberg: Springer-Verlag 2011.

5. Lacey JV, Jr., Chia VM, Rush BB, Carreon DJ, Richesson DA, Ioffe OB, Ronnett BM, Chatterjee N, Langholz B, Sherman ME, Glass AG. Incidence rates of endometrial hyperplasia, endometrial cancer and hysterectomy from 1980 to 2003 within a large prepaid health plan. *Int J Cancer*. 2012;131(8):1921-9. doi:10.1002/ijc.27457.

6. Amant F, Moerman P, Neven P, Timmerman D, Van Limbergen E, Vergote I. Endometrial cancer. *The Lancet*. 2005;366(9484):491-505.

7. World Cancer Research Fund / American Institute for Cancer Research. *Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global perspective*. Washington DC: AICR; 2007.

8. World Cancer Research Fund/American Institute for Cancer Research. *Continuous Update Project Report. Food, Nutrition, Physical Activity, and the prevention of Endometrial Cancer*. Available at <http://www.dietandcancerreport.org>. 2013.

9. Lindemann K, Eskild A, Vatten LJ, Bray F. Endometrial cancer incidence trends in Norway during 1953–2007 and predictions for 2008–2027. *International Journal of Cancer*. 2010;127(11):2661-8. doi:10.1002/ijc.25267.

10. Moore SC, Lee I, Weiderpass E, Campbell PT, Sampson JN, Kitahara CM, Keadle SK, Arem H, Gonzalez Abd, Hartge P, Adami H-O, Blair CK, Borch KB, Boyd E, Check DP, Fournier A, Freedman ND, Gunter M, Johansson M, Khaw K-T, Linet MS, Orsini N, Park Y, Riboli E, Robien K, Schairer C, Sesso H, Spriggs M, Dusen RV, Wolk A, Matthews CE, Patel AV. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA internal medicine*. 2016;176(6):816-25. doi:10.1001/jamainternmed.2016.1548.

11. Conroy MB, Sattelmair JR, Cook NR, Manson JE, Buring JE, Lee I-M. Physical activity, adiposity, and risk of endometrial cancer. *Cancer Causes & Control*. 2009;20(7):1107-15. doi:10.1007/s10552-009-9313-3.

12. Friedenreich C, Cust A, Lahmann PH, Steindorf K, Boutron-Ruault M-C, Clavel-Chapelon F, Mesrine S, Linseisen J, Rohrmann S, Pischon T, Schulz M, Tjønneland A, Johnsen NF, Overvad K, Mendez M, Arguelles MV, Garcia CM, Larrañaga N, Chirlaque M-D, Ardanaz E, Bingham S, Khaw K-T, Allen N, Key T, Trichopoulou A, Dilis V, Trichopoulos D, Pala V, Palli D, Tumino R, Panico S, Vineis P, Bueno-de-Mesquita HB, Peeters PHM, Monninkhof E, Berglund G, Manjer J, Slimani N, Ferrari P, Kaaks R, Riboli E. Physical activity and risk of endometrial cancer: The European prospective investigation into cancer and nutrition. *International Journal of Cancer*. 2007;121(2):347-55. doi:10.1002/ijc.22676.

13. Dieli-Conwright CM, Ma H, Lacey JV, Henderson KD, Neuhausen S, Horn-Ross PL, Deapen D, Sullivan-Halley J, Bernstein L. Long-term and baseline recreational physical activity and risk of endometrial cancer: the California Teachers Study. *Br J Cancer*. 2013;109(3):761-8. doi:10.1038/bjc.2013.61.

14. Land SR, Liu Q, Wickerham DL, Costantino JP, Ganz PA. Cigarette Smoking, Physical Activity, and Alcohol Consumption as Predictors of Cancer Incidence among Women at High Risk of Breast Cancer in the NSABP P-1 Trial. *Cancer Epidemiology Biomarkers & Prevention*. 2014;23(5):823-32. doi:10.1158/1055-9965.epi-13-1105-t.

15. Du M, Kraft P, Eliassen AH, Giovannucci E, Hankinson SE, De Vivo I. Physical activity and risk of endometrial adenocarcinoma in the Nurses' Health Study. *International Journal of Cancer*. 2014;134(11):2707-16. doi:10.1002/ijc.28599.
16. Friberg E, Mantzoros CS, Wolk A. Physical Activity and Risk of Endometrial Cancer: A Population-Based Prospective Cohort Study. *Cancer Epidemiology Biomarkers & Prevention*. 2006;15(11):2136-40. doi:10.1158/1055-9965.epi-06-0465.
17. Schouten LJ, Goldbohm RA, van den Brandt PA. Anthropometry, Physical Activity, and Endometrial Cancer Risk: Results From The Netherlands Cohort Study. *Journal of the National Cancer Institute*. 2004;96(21):1635-8. doi:10.1093/jnci/djh291.
18. Murali R, Soslow RA, Weigelt B. Classification of endometrial carcinoma: more than two types. *The Lancet Oncology*. 2014;15(7):e268-e78. doi:[http://dx.doi.org/10.1016/S1470-2045\(13\)70591-6](http://dx.doi.org/10.1016/S1470-2045(13)70591-6).
19. Lund E, Kumle M, Braaten T, Hjartåker A, Bakken K, Eggen A, Gram I. External validity in a population-based national prospective study – the Norwegian Women and Cancer Study (NOWAC). *Cancer Causes and Control*. 2003;14(10):1001-8. doi:10.1023/B:CACO.0000007982.18311.2e.
20. Lund E, Dumeaux V, Braaten T, Hjartåker A, Engeset D, Skeie G, Kumle M. Cohort Profile: The Norwegian Women and Cancer Study (NOWAC) Kvinner og kreft. *Int J Epidemiol*. 2008;37(1):36-41. doi:10.1093/ije/dym137.
21. Borch KB, Ekelund U, Brage S, Lund E. Criterion validity of a 10-category scale for ranking physical activity in Norwegian women. *Int J Behav Nutr Phys Act*. 2012;9(1):2.
22. Collaborators MWS. Breast cancer and hormone-replacement therapy in the Million Women Study. *The Lancet*. 2003;362(9382):419-27. doi:[http://dx.doi.org/10.1016/S0140-6736\(03\)14065-2](http://dx.doi.org/10.1016/S0140-6736(03)14065-2).
23. Waaseth M, Bakken K, Dumeaux V, Olsen K, Rylander C, Figenschau Y, Lund E. Hormone replacement therapy use and plasma levels of sex hormones in the Norwegian Women and Cancer Postgenome Cohort - a cross-sectional analysis. *BMC Women's Health*. 2008;8(1):1.
24. WHO. International Statistical Classification of Diseases and Related Health Problems 10th Revision. <http://apps.who.int/classifications/icd10/browse/2015/en>. 2010. Accessed 27.11.2014.
25. Hu P, Tsiatis AA, Davidian M. Estimating the Parameters in the Cox Model When Covariate Variables are Measured with Error. *Biometrics*. 1998;54(4):1407-19. doi:10.2307/2533667.
26. White IR, Carlin JB. Bias and efficiency of multiple imputation compared with complete-case analysis for missing covariate values. *Statistics in Medicine*. 2010;29(28):2920-31. doi:10.1002/sim.3944.
27. Sterne J, White I, Carlin J, Spratt M, Royston P, Kenward M, Wood A, Carpenter, Jr. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. 2009. doi:10.1136/bmj.b2393.
28. White IR, Royston P. Imputing missing covariate values for the Cox model. *Statistics in Medicine*. 2009;28(15):1982-98. doi:10.1002/sim.3618.
29. Rubin DB. Multiple Imputation After 18+ Years. *Journal of the American Statistical Association*. 1996;91(434):473-89. doi:10.2307/2291635.
30. Benedict N. PUNAF: Stata module to compute population attributable fractions for cohort studies. <https://ideas.repec.org/c/boc/bocode/s457193.html>. 2010. Accessed 23 June 2016.
31. Colbert LH, Lacey JV, Schairer C, Albert P, Schatzkin A, Albanes D. Physical activity and risk of endometrial cancer in a prospective cohort study (United States). *Cancer Causes & Control*. 2003;14(6):559-67. doi:10.1023/a:1024866827775.
32. Schouten LJ, Goldbohm RA, van den Brandt PA. Anthropometry, Physical Activity, and Endometrial Cancer Risk: Results From The Netherlands Cohort Study. *International Journal of Gynecological Cancer*. 2006;16 (suppl. 2):492.
33. Gierach GL, Chang S-C, Brinton LA, Lacey JV, Hollenbeck AR, Schatzkin A, Leitzmann MF. Physical activity, sedentary behavior, and endometrial cancer risk in the NIH-AARP Diet and Health Study. *International Journal of Cancer*. 2009;124(9):2139-47. doi:10.1002/ijc.24059.

1
2
3
4
5
6
7 521 34. Terry P, Baron JA, Weiderpass E, Yuen J, Lichtenstein P, Nyrén O. Lifestyle and endometrial cancer
8 522 risk: a cohort study from the Swedish twin registry. *International Journal of Cancer*. 1999;82(1):38-
9 523 42. doi:10.1002/(SICI)1097-0215(19990702)82:1<38::AID-IJC8>3.0.CO;2-Q.
10 524 35. Folsom AR, Demissie Z, Harnack L. Glycemic index, glycemic load, and incidence of endometrial
11 525 cancer: the Iowa Women's Health Study. *Nutrition & Cancer*. 2003;46(2):119-24 6p.
12 526 36. Furberg A-S, Thune I. Metabolic abnormalities (hypertension, hyperglycemia and overweight),
13 527 lifestyle (high energy intake and physical inactivity) and endometrial cancer risk in a Norwegian
14 528 cohort. *International Journal of Cancer*. 2003;104(6):669-76. doi:10.1002/ijc.10974.
15 529 37. Patel AV, Feigelson HS, Talbot JT, McCullough ML, Rodriguez C, Patel RC, Thun MJ, Calle EE. The
16 530 role of body weight in the relationship between physical activity and endometrial cancer: Results
17 531 from a large cohort of US women. *International Journal of Cancer*. 2008;123(8):1877-82.
18 532 doi:10.1002/ijc.23716.
19 533 38. Moradi T, Nyrén O, Bergström R, Gridley G, Linet M, Wolk A, Dosemeci M, Adami H-O. Risk for
20 534 endometrial cancer in relation to occupational physical activity: A nationwide cohort study in
21 535 Sweden. *International Journal of Cancer*. 1998;76(5):665-70. doi:10.1002/(SICI)1097-
22 536 0215(19980529)76:5<665::AID-IJC9>3.0.CO;2-O.
23 537 39. Weiderpass E, Pukkala E, Vasama-Neuvonen K, Kauppinen T, Vainio H, Paakkulainen H, Boffetta P,
24 538 Partanen T. Occupational exposures and cancers of the endometrium and cervix uteri in Finland.
25 539 *American Journal of Industrial Medicine*. 2001;39(6):572-80. doi:10.1002/ajim.1056.
26 540 40. Friedenreich CM. Physical Activity and Cancer Prevention: From Observational to Intervention
27 541 Research. *Cancer Epidemiology Biomarkers & Prevention*. 2001;10(4):287-301.
28 542 41. Cust AE. Physical Activity and Gynecologic Cancer Prevention. Physical Activity and Cancer. In:
29 543 Courneya KS, Friedenreich CM, editors. *Recent Results in Cancer Research*. Berlin Heidelberg:
30 544 Springer; 2011. p. 159-85.
31 545 42. Mathers C, Stevens GA, Mascarenhas M. Global Health Risks: Mortality burden of disease
32 546 attributable to selected major risk. Switzerland: WHO2009. Report No.: 9789241563871.
33 547 43. Parkin DM. 9. Cancers attributable to inadequate physical exercise in the UK in 2010. *Br J Cancer*.
34 548 2011;105(S2):S38-S41.
35 549 44. Olsen CM, Wilson LF, Nagle CM, Kendall BJ, Bain CJ, Pandeya N, Webb PM, Whiteman DC.
36 550 Cancers in Australia in 2010 attributable to insufficient physical activity. *Australian and New Zealand*
37 551 *Journal of Public Health*. 2015;39(5):458-63. doi:10.1111/1753-6405.12469.
38 552 45. World Cancer Research Fund/American Institute for Cancer Research. Policy and Action for
39 553 Cancer Prevention. Food, Nutrition and Physical Activity: A Global Perspective. AICR: Washington, DC.
40 554 2009.
41 555 46. Borch KB, Braaten T, Lund E, Weiderpass E. Physical activity and mortality among Norwegian
42 556 women - the Norwegian Women and Cancer Study. *Clin Epidemiol*. 2011;3(1):229-35.
43 557 doi:10.1186/1479-5868-9-2.
44 558 47. Skeie G, Mode N, Henningsen M, Borch KB. Validity of self-reported body mass index among
45 559 middle-aged participants in the Norwegian Women and Cancer study. *Clinical Epidemiology*.
46 560 2015;7:313-23. doi:10.2147/CLEP.S83839.
47 561 48. Hansen BH, Anderssen SA, Steene-Johannessen J, Ekelund U, Nilsen AK, Andersen ID, Dalene KE,
48 562 Kolle E. Fysisk aktivitet og sedat tid blant voksne og eldre i Norge-Nasjonal kartlegging 2014-2015.
49 563 Helsedirektoratet, Oslo2015.
50
51
52
53
54
55
56
57
58
59
60

Formatted: Font: Calibri, Norwegian (Bokmål), Do not check spelling or grammar

Formatted: Norwegian (Bokmål)

Formatted: Font: Calibri, Norwegian (Bokmål), Do not check spelling or grammar

Table 1 Selected baseline characteristics of participants in the Norwegian Women and Cancer Study by physical activity level (n=82,759)

Characteristic	Physical activity level				
	1 to 4 N=21,953 (26.5%)		5 to 10 N=60,806 (73.5%)		
	1 to 2 N=3855 (4.7%)	3 to 4 N=18,098 (21.9%)	5 to 6 N=35,551 (43.0%)	7 to 8 N=20,991 (25.4%)	9 to 10 N=4264 (5.2%)
Age (mean, \pm SE)	53.2 (.11)	52.1 (.05)	51.5 (.03)	51.3 (.04)	51.9 (.10)
Person-years at risk ¹	48 995	236 571	462 837	266 512	54 318
Overall endometrial cancer (total cases n=687)	56 (8.2%)	181 (26.3%)	283 (41.2%)	139 (20.2%)	28 (4.1%)
Endometrioid (total cases n=473)	41 (8.7%)	134 (28.3%)	185 (39.1%)	100 (21.1%)	13 (2.7%)
Type 1 endometrial cancer ² (total cases n=576)	47 (8.2%)	161 (28.0%)	226 (39.2%)	122 (21.2%)	20 (3.5%)
Other subtypes ³ (total cases n=111)	9 (8.1%)	20 (18%)	57 (51.4%)	17 (15.3%)	8 (7.2%)
BMI (mean, \pm SE)	26.9 (.09)	25.8 (.03)	24.6 (.02)	23.8 (.02)	23.6 (.05)
Missing (%)	2.2	2.0	1.7	1.7	2.2
Age at menarche (mean, \pm SE)	13.2 (.03)	13.3 (.01)	13.3 (.007)	13.7 (.01)	13.4 (.02)
Missing (%)	1.6	1.6	1.4	1.2	1.7
Parity (%)					
Nulliparity	10.4	9.2	8.2	8.0	8.2
1-2	51.6	52.7	53.2	53.0	50.4
≥ 3	38.0	38.1	38.5	39.0	41.4
Ever use of oral contraceptives (%)	49.6	53.9	53.8	54.8	50.4
Missing (%)	3.9	2.9	2.8	3.2	4.3
Menopausal status (%)					
Premenopausal	38.4	47.4	50.0	48.8	44.4
Perimenopause	7.0	5.9	5.9	6.5	6.1
Postmenopausal	49.0	41.9	39.0	39.4	44.4
Hormonal therapy use <53 years	3.0	2.8	2.8	3.1	3.3
Missing	2.6	2.0	2.3	2.3	1.8
Ever use of hormonal therapy (%)	36.9	35.5	32.5	31.4	29.6
Missing (%)	2.8	2.4	2.6	2.4	3.4
Years of education (mean, \pm SE)	11.5 (.06)	12.2 (.03)	12.3 (.02)	12.6 (.02)	11.9 (.06)

Missing (%)	6.6	5.2	4.5	4.9	6.0
Smoking status (%)					
Never	29.6	36.1	37.3	37.4	35.2
Former	29.8	31.2	32.6	35.0	33.3
Current	39.1	31.4	28.9	26.3	30.0
Missing (%)	1.5	1.3	1.2	1.4	1.5
Alcohol consumption, mean ±SE (grams/day)	3.0 (.03)	3.4 (.03)	3.4 (.02)	3.5 (.03)	3.1 (.06)
Missing (%)	2.3	1.7	1.7	1.8	2.4
Diabetes mellitus (%)	4.4	2.1	1.6	1.3	1.6
Missing (%)	27.3	20.7	17.3	15.3	17.1

¹Total person years=1,069,232; average follow-up time 12.92 years (SD=3.65)

²Type 1 includes adenocarcinoma, endometrioid, squamous types

³Other subtypes are cases not equal to Type 1

Table 2 Relative risk estimates of endometrial cancer (overall, type 1, and endometrioid) by physical activity (PA) level at baseline and follow-up in the Norwegian Women and Cancer Study (n=82 759)

	Endometrial cancer	Models ^{1, 2, 3, 4}	1 to 2	3 to 4	5 to 6	7 to 8	9 to 10	p_trend
Baseline	Overall endometrial cancer	Crude n=687	1.73 (1.30-2.31)	1.21 (1.00-1.46)	1.00	0.87 (0.71-1.07)	0.83 (0.56-1.22)	0.00
		PA + BMI n= 673	1.47 (1.09-1.97)	1.09 (0.9-1.32)	1.00	0.91 (0.74-1.11)	0.86 (0.58-1.27)	0.01
		Multivariable without BMI n=607	1.85 (1.36-2.53)	1.28 (1.05-1.56)	1.00	0.89 (0.72-1.11)	0.71 (0.45-1.12)	0.00
		Multivariable n=593	1.60 (1.16-2.20)	1.15 (0.94-1.41)	1.00	0.92 (0.74-1.15)	0.73 (0.45-1.16)	0.01
	Type 1 endometrial cancer	Crude n=576	1.83 (1.33-2.51)	1.35 (1.10-1.65)	1.00	0.96 (0.77-1.19)	0.74 (0.47-1.17)	0.00
		PA + BMI n=564	1.51 (1.10-2.09)	1.20 (0.97-1.47)	1.00	1.01 (0.81-1.26)	0.76 (0.48-1.22)	0.01
		Multivariable without BMI n=510	1.97 (1.4-2.78)	1.47 (1.19-1.82)	1.00	1.00 (0.79-1.26)	0.68 (0.4-1.15)	0.00
		Multivariable n=498	1.66 (1.17-2.36)	1.31 (1.05-1.63)	1.00	1.04 (0.82-1.32)	0.69 (0.40-1.18)	0.00
	Endometrioid subtype	Crude n=473	1.96 (1.39-2.75)	1.37 (1.10-1.72)	1.00	0.96 (0.75-1.22)	0.59 (0.33-1.03)	0.00
		PA + BMI n=463	1.59 (1.12-2.25)	1.20 (0.96-1.51)	1.00	1.00 (0.79-1.29)	0.59 (0.33-1.06)	0.004
		Multivariable without BMI n=428	2.25 (1.57-3.22)	1.51 (1.2-1.91)	1.00	1.01 (0.78-1.3)	0.61 (0.33-1.13)	0.00
		Multivariable n=418	1.83 (1.26-2.64)	1.32 (1.04-1.67)	1.00	1.06 (0.82-1.37)	0.60 (0.32-1.15)	0.001
Repeated measurements PA	Overall	Crude n= 450	1.78 (1.22-2.58)	1.31 (1.04-1.65)	1.00	0.86 (0.67-1.10)	0.87 (0.54-1.40)	0.00
		PA + BMI n=438	1.48 (1.01-2.17)	1.18 (0.93-1.49)		0.91 (0.71-1.17)	0.98 (0.61-1.57)	0.04
		Multivariable without BMI n=393	1.80 (1.19-2.72)	1.35 (1.06-1.73)	1.00	0.92 (0.71-1.2)	0.71 (0.4-1.26)	0.00
		Multivariable n=381	1.54 (1.01-2.35)	1.22 (0.95-1.58)	1.00	0.97 (0.74-1.26)	0.80 (0.45-1.41)	0.02
	Type 1	Crude n = 387	1.98 (1.34-2.91)	1.39 (1.09-1.78)	1.00	0.83 (0.63-1.09)	0.93 (0.56-1.53)	0.00
		PA + BMI n=377	1.60 (1.08-2.39)	1.25 (0.97-1.61)	1.00	0.88 (0.67-1.17)	1.06 (0.64-1.75)	0.01

		Multivariable without BMI n=337	2.10 (1.38-3.21)	1.46 (1.12-1.9)	1.00	0.94 (0.71-1.25)	0.8 (0.44-1.45)	0.00
		Multivariable n=327	1.75 (1.13-2.71)	1.33 (1.01-1.74)	1.00	0.99 (0.74-1.33)	0.91 (0.5-1.65)	0.01
	Endometrioid	Crude n = 321	1.90 (1.24-2.89)	1.30 (1.00-1.70)	1.00	0.72 (0.53-0.97)	0.61 (0.32-1.16)	0.00
		PA + BMI n=311	1.55 (1.00-2.39)	1.18 (0.9-1.55)	1.00	0.76 (0.55-1.03)	0.7 (0.37-1.34)	0.01
		Multivariable without BMI n=290	2.15 (1.38-3.34)	1.34 (1.01-1.79)	1.00	0.78 (0.57-1.07)	0.65 (0.33-1.28)	0.00
		Multivariable n=280	1.77 (1.12-2.81)	1.22 (0.91-1.63)	1.00	0.82 (0.60-1.14)	0.75 (0.38-1.49)	0.01

¹ Crude model with age as time-variable.
² Model with adjustment for BMI.
³ Multivariable model adjusted for use of hormone therapy, oral contraceptive use, years of education, smoking, alcohol consumption (gram per day).
⁴ Multivariable model same as above, but with adjustments for BMI included.
BMI: body mass index

Table 3 Relative risk estimates of overall endometrial cancer by physical activity (PA) level at baseline and follow-up according to body mass index (BMI) status at baseline in the Norwegian Women and Cancer study (n=82,759)

Complete case models	Model ^{1,2}	BMI	1 to 2	3 to 4	5 to 6	7 to 10	p_trend
Baseline	Crude	Normal BMI n=318	1.28 (0.74-2.21)	1.06 (0.79-1.42)	1.00	0.83 (0.64-1.07)	0.10
		Overweight n=234	1.04 (0.60-1.83)	1.02 (0.75-1.39)	1.00	0.90 (0.64-1.26)	0.66
		Obese n=121	2.50 (1.49-4.20)	1.45 (0.92-2.29)	1.00	1.32 (0.74-2.36)	0.01
	Multivariable	Normal BMI n=281	1.32 (0.71-2.45)	1.19 (0.87-1.62)	1.00	0.86 (0.66-1.14)	0.07
		Overweight n=212	0.95 (0.51-1.77)	1.00 (0.73-1.39)	1.00	0.90 (0.63-1.28)	0.76
		Obese n=100	3.08 (1.76-5.39)	1.55 (0.93-2.56)	1.00	0.98 (0.48-1.99)	0.00
Repeated measurements PA	Crude	Normal BMI n=182	1.26 (0.58-2.73)	0.85 (0.56-1.30)	1.00	0.80 (0.58-1.12)	0.39
		Overweight n=169	1.44 (0.74-2.82)	1.39 (0.97-2.01)	1.00	1.16 (0.78-1.71)	0.61
		Obese n=87	1.78 (0.94-3.37)	1.39 (0.84-2.30)	1.00	0.73 (0.34-1.56)	0.01
	Multivariable	Normal BMI n=164	1.17 (0.47-2.91)	0.90 (0.58-1.41)	1.00	0.86 (0.61-1.21)	0.53
		Overweight n=147	1.15 (0.52-2.53)	1.35 (0.92-2.00)	1.00	1.17 (0.78-1.77)	0.81
		Obese n=70	2.45 (1.22-4.91)	1.63 (0.92-2.89)	1.00	0.56 (0.21-1.50)	0.00

¹ Crude model with age as time-variable.

² Multivariable model adjusted for use of hormone therapy, oral contraceptive use, years of education, smoking, alcohol consumption (gram/day).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 4: Relative risk estimates of endometrial cancer (overall, type 1 and endometrioid) by physical activity (PA) level at baseline and follow-up with imputation for missing variables in the Norwegian Women and Cancer study (n=82,759)

Imputation	Models ^{1,2,3,4}	1 to 2	3 to 4	5 to 6	7 to 8	9 to 10	p_trend
Overall	Crude n=687	1.76 (1.27-2.44)	1.34 (1.09-1.65)	1.00	0.80 (0.65-1.00)	0.73 (0.48-1.12)	0.00
	PA + BMI n= 687	1.47 (1.05-2.05)	1.22 (0.98-1.50)	1.00	0.85 (0.69-1.06)	0.80 (0.52-1.22)	0.001
	Multivariable without BMI n=687	1.87 (1.32-2.63)	1.40 (1.13-1.74)	1.00	0.85 (0.69-1.07)	0.71 (0.45-1.13)	0.02
	Multivariable n=687	1.58 (1.11-2.24)	1.28 (1.02-1.59)	1.00	0.90 (0.72-1.13)	0.78 (0.49-1.22)	0.02
Type 1	Crude n=576	1.87 (1.23-2.83)	1.30 (1.00-1.70)	1.00	0.71 (0.53-0.96)	0.60 (0.31-1.14)	0.00
	PA + BMI n= 576	1.51 (0.99-2.31)	1.16 (0.89-1.52)	1.00	0.77 (0.57-1.04)	0.67 (0.35-1.27)	0.001
	Multivariable without BMI n=576	2.05 (1.34-3.15)	1.34 (1.02-1.77)	1.00	0.77 (0.57-1.05)	0.66 (0.35-1.27)	0.00
	Multivariable n=576	1.66 (1.08-2.57)	1.19 (0.90-1.58)	1.00	0.83 (0.61-1.13)	0.74 (0.39-1.42)	0.002
Endometrioid	Crude n=473	1.89 (1.28-2.79)	1.45 (1.13-1.85)	1.00	0.76 (0.58-0.99)	0.48 (0.24-0.96)	0.00
	PA + BMI n= 473	1.51 (1.01-2.26)	1.28 (1.00-1.64)	1.00	0.82 (0.63-1.07)	0.54 (0.27-1.08)	0.01
	Multivariable without BMI n=473	2.11 (1.42-3.14)	1.53 (1.18-1.97)	1.00	0.83 (0.63-1.09)	0.54 (0.27-1.07)	0.00
	Multivariable n=473	1.70 (1.13-2.56)	1.35 (1.05-1.75)	1.00	0.89 (0.68-1.17)	0.60 (0.30-1.20)	0.00

¹ Crude model with age as time-variable.

² Model with adjustment for BMI.

³ Multivariable model adjusted for use of hormone therapy, oral contraceptive use, years of education, smoking, alcohol consumption (gram per day).

⁴ Multivariable model same as above, but with adjustment for BMI included.

BMI: body mass index

Table 5: Population attributable fraction (PAF) for the proportion of endometrial cancer in the population that would be avoided if low physical activity (PA) level increased

		Overall					
		Model 1 ¹ n= 673			Model 2 ² n= 593		
PA level	Pe	PAF	PAF 95% CI	RR= 95% CI	PAF	PAF 95% CI	RR= 95% CI
[1-4] to [5-10]	26.5 %	17%	[2.3-29.5]	1.21 [1.02-1.42]	21.9%	[7.1-34.3]	1.28 [1.08-1.52]

¹Model 1: measured at baseline with age as time variable and adjusted for BMI.

²Model 2: Measured at baseline in a multivariable model adjusted for BMI, oral contraceptive use, hormone therapy use, years of education, smoking status, and alcohol consumption (grams per day).

Pe: proportion of low physical activity levels, RR: relative risk, CI: confidence interval